

LETTER • OPEN ACCESS

## Greenhouse gas emissions from windrow composting of organic wastes: Patterns and emissions factors

To cite this article: Sintana E Vergara and Whendee L Silver 2019 *Environ. Res. Lett.* **14** 124027

View the [article online](#) for updates and enhancements.



## LETTER

## Greenhouse gas emissions from windrow composting of organic wastes: Patterns and emissions factors

## OPEN ACCESS

RECEIVED  
6 August 2019REVISED  
25 October 2019ACCEPTED FOR PUBLICATION  
29 October 2019PUBLISHED  
4 December 2019

Original content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](#).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

Sintana E Vergara<sup>1</sup> and Whendee L Silver

Department of Environmental Science, Policy, and Management, University of California, Berkeley, Berkeley, CA, United States of America  
<sup>1</sup> Present address: Department of Environmental Resources Engineering, Humboldt State University, 1 Harpst Street, Arcata CA 95521, United States of America.

E-mail: [vergara@humboldt.edu](mailto:vergara@humboldt.edu)**Keywords:** compost, biogeochemistry, micrometeorological mass balance, lifecycle assessment, controls on decompositionSupplementary material for this article is available [online](#)**Abstract**

Direct emissions from commercial-scale composting are uncertain. We used micrometeorological methods to continuously measure greenhouse gas (CO<sub>2</sub>, CH<sub>4</sub>, N<sub>2</sub>O) emissions from full composting of green waste and manure. We measured oxygen (O<sub>2</sub>), moisture, and temperature continuously inside the composting pile, and analyzed chemical and physical characteristics of the feedstock weekly as potential drivers of emissions. Temperature, moisture, and O<sub>2</sub> all varied significantly by week. Feedstock porosity, C:N, and potential N mineralization all declined significantly over time. Potential net nitrification remained near zero throughout. CH<sub>4</sub> and CO<sub>2</sub> fluxes, indicators of feedstock lability, were variable, and most emissions (75% and 50% respectively) occurred during the first three weeks of composting. Total CH<sub>4</sub> emitted was  $1.7 \pm 0.32$  g CH<sub>4</sub> kg<sup>-1</sup> feedstock, near the median literature value using different approaches (1.4 g CH<sub>4</sub> kg<sup>-1</sup>). N<sub>2</sub>O concentrations remained below the instrument detection. Oxygen, moisture and temperature exhibited threshold effects on CH<sub>4</sub> emissions. Net lifecycle emissions were negative ( $-690$  g CO<sub>2</sub>-e kg<sup>-1</sup>), however, after considering avoided emissions and sinks. Managing composting piles to minimize methanogenesis—by maintaining sufficient O<sub>2</sub> concentrations, and focusing on the first three weeks—could reduce emissions, contributing to the climate change mitigation benefit of composting.

**Introduction**

Organic waste in landfills and manure slurries is a large source of greenhouse gas emissions globally [1, 2]. Manure alone accounts for 10% of global agricultural emissions [2]. Diverting organic waste to composting may lower greenhouse gas emissions, but there is uncertainty regarding emissions from the composting process. Composting, the aerobic degradation of organic matter mediated by microbes, is used to manage wastes and recycle nutrients into the soil [3]. Under aerobic conditions, organic waste is converted to stable material; microbial respiration emits mostly carbon dioxide (CO<sub>2</sub>) and N mineralization predominantly produces ammonium (NH<sub>4</sub><sup>+</sup>). Compost piles often have pockets of anaerobiosis, and even well-managed composting emits methane (CH<sub>4</sub>) and

nitrous oxide (N<sub>2</sub>O) [1], greenhouse gases that trap 34 and 298 times more heat than CO<sub>2</sub>, over 100 years [4].

Aerobic composting follows four phases. In the first mesophilic (25 °C–40 °C) stage, primary decomposers break down easily degradable material, releasing CO<sub>2</sub> and heat. High redox potentials allow for the presence of nitrifying and denitrifying bacteria, potentially emitting N<sub>2</sub>O [5]. The thermophilic (40 °C–65 °C) stage features increasing microbial activity and temperatures, which can favor oxygen (O<sub>2</sub>) consumption and methanogenesis [5]. The second mesophilic stage begins as easily degradable material is consumed, and thermophilic activity and compost temperature decrease. During maturation, bacterial numbers decline and fungal populations increase as easily degraded material is exhausted. Recalcitrant material dominates and temperature declines to ambient levels [6].

Emissions from composting at a commercial scale are not well characterized. Of the studies that have measured emissions from composting e.g. [7–10], few measured continuously, few measured at field scale [10] and estimates of total emissions vary widely (table S1 is available online at [stacks.iop.org/ERL/14/124027/mmedia](https://stacks.iop.org/ERL/14/124027/mmedia)). One reason composting emissions are poorly constrained is that effective measurement requires technologically advanced instrumentation due to the scale and heterogeneity of compost piles. Methods used to measure greenhouse gases from composting include micrometeorological (e.g. eddy covariance, mass balance) [11] and non-micrometeorological techniques (e.g. static chambers). Chamber techniques measure at small scales (<1 m) and introduce sources of bias, including altering pressure and concentration gradients, physically disturbing the source, leakage, difficulty in capturing spatial and temporal variation, and potential for significant human error [9, 12, 13]. Enclosing the emitting source [7] facilitates emission measurement, but introduces artifacts from altering ambient conditions and degradation dynamics.

Micrometeorological methods use physical, chemical, and engineering principles to estimate greenhouse gas fluxes by continuously measuring gas concentrations and wind dynamics. The micrometeorological mass balance (MMB) approach measures gas concentrations going into and coming out of a control volume surrounding an emitting source. The source's emission rate is calculated by subtracting the input from the output flux [11].

Three previous studies [7, 8, 12] used micrometeorological approaches to measure emissions from composting, though none measured fluxes from a commercial process, nor from composting manure with green waste. These studies provided a basis upon which the methodology was developed. Amlinger *et al* (2008) used a large chamber enclosing a compost windrow and measured gas concentrations at the inflows and outflows. This approach captured emissions, though covering the pile may have altered gas fluxes by increasing temperature and moisture, and affecting the concentration gradient from the source to the atmosphere. Sommer *et al* (2004) and Kent (2010) used a MMB approach. Sommer (2004) measured concentrations of gases upstream and downstream of a circular manure pile, not the windrow formation typical of commercial composting operations, and Kent (2010) analyzed emissions from green waste composting in windrows, but the study encountered instrumentation malfunctions, and was short in duration (5 weeks).

Understanding how environmental variables influence greenhouse gas fluxes is critical for modeling, extrapolating results, and developing strategies to reduce emissions. Though composting is mostly aerobic, the heterogeneity of the feedstock, temperature and moisture, microbial activity, and the pile structure

can create variable redox conditions. In the absence of oxygen ( $O_2$ ), a succession of microbes converts carbohydrates in the organic waste to  $CO_2$  and  $CH_4$  [14].  $O_2$  availability is affected by feedstock porosity [7, 15], turning frequency [5], and pile size. Once  $CH_4$  is produced, it may be emitted to the atmosphere or oxidized to  $CO_2$  within the pile. The balance between  $CH_4$  production and oxidation is likely controlled by redox potential [16] and is affected by temperature and moisture, which control  $O_2$  solubility and biological activity [17, 18].

Nitrification, the conversion of  $NH_4^+$  to  $NO_3^-$ , and denitrification, the conversion of  $NO_3^-$  to nitrogen gas ( $N_2$  and  $N_2O$ ), are the major pathways leading to  $N_2O$  production and consumption [19]. Net emission of  $N_2O$  is dependent on the controls on both processes. The production of  $NO_3^-$  during composting poses a potential water eutrophication threat. Nitrification is regulated by  $NH_4^+$  supply, pH, and redox potential; denitrification is dependent on  $NO_3^-$  supply, C availability, and redox.  $N_2O$  fluxes are expected to increase as the C:N ratio of organic matter decreases, as the N availability increases [20] and as  $O_2$  concentrations decline [19].

This study is the first to continuously measure greenhouse gas emissions from the commercial-scale composting of manure and green waste. Three questions guided our research: (1) how do environmental and biogeochemical characteristics vary during composting?; (2) what are the whole-pile emissions of  $CO_2$ ,  $CH_4$ , and  $N_2O$ , and how are they related to environmental and biogeochemical variables?; and (3) what are the net lifecycle greenhouse gas emissions from composting green waste and manure? We used field experiments, laboratory assays, and lifecycle modeling to answer these questions.

## Methods

### Site description and experimental set-up

The experiment was conducted at the West Marin Composting Facility in Nicasio, California, from February through September 2016. A windrow pile (15 m × 2 m × 4 m) comprised of manure (mostly cattle, with goat, horse, chicken) and yard waste (branches, leaves, grass) was composted. The starting pile was 55% manure by volume and 91% manure by mass. The pile was managed as a commercially-produced compost pile: with weekly turning using a large-scale mechanical windrow turner (Scarab 18), periodic watering, and a 98 d duration.

### Sensor system and laboratory analyses

Conditions inside the piles were monitored continuously using 27 automated sensors. Nine  $O_2$  (Apogee SO-110), temperature (Campbell 107), and moisture (Campbell CS616) sensors each were placed in three transects along the length of the pile, at three heights

(0.50 m, 1 m, 1.5 m), in the center of the pile (figures S1-2). Sensors were removed briefly for weekly pile turning (<60 min). Compost grab samples were collected weekly into 1 gallon Ziploc bags from three heights (0.5 m, 1 m, 1.5 m—at 2 m horizontal depth) before and after pile turning ( $n = 6$ ). Samples were analyzed for gravimetric moisture content by weighing 10 g samples before and after drying at 105° C for 24 h. Compost pH was measured by suspending 3 g of sample in 12 g of water [21]. We estimated porosity by weighing a mason jar with a volume of 100 ml of collected compost, adding 100 ml distilled water, and re-weighing to determine the volume of the pore space [22] ( $n = 5$  per sample). Note that collecting samples for porosity may result in disturbance to the sample and thus not represent exact in situ conditions. Potential net N mineralization and nitrification were determined using dark laboratory incubations; 3.5 g compost samples were extracted before and after incubation (7 d) in 75 ml of 2 M KCl ( $n = 3$  per sample) [23]. Total C and N were measured on air-dried, sieved (2 mm) and ground samples using an elemental analyzer (Carlo Erba Elantech, Lakewood, NJ, USA).

### MMB method

The pile was outfitted with a greenhouse gas and wind measurement system. Four gas towers (figure S2, A–D) were placed at cardinal directions along each edge of the windrow, 1 m from the pile edge; each of these towers held four gas intakes (heights of 0.25, 1, 2, and 3.5 m). Air samples were continuously drawn through 16 Teflon tubes, delivering gas samples to a G2308 cavity ring down laser spectrometer (Picarro, Santa Clara, CA), which measured real-time CO<sub>2</sub>, N<sub>2</sub>O and CH<sub>4</sub> concentrations. The instrument was calibrated using a three-point calibration curve, using zero (pure N<sub>2</sub> gas), intermediate (304 ppm CO<sub>2</sub>, 0.513 ppm N<sub>2</sub>O, 1.05 ppm CH<sub>4</sub>), and high (1000 ppm CO<sub>2</sub>, 10 ppm N<sub>2</sub>O, 10 ppm CH<sub>4</sub>) standard concentrations. The instrument sensitivity is high for CH<sub>4</sub> and N<sub>2</sub>O (raw precision <10 ppb and <25 ppb, respectively), but low for CO<sub>2</sub> (<20 ppm). While one intake port delivered gases to the greenhouse gas analyzer (1 min per intake), the other 15 lines were flushed with ambient air.

Two wind towers on the NW and SE corners of the windrow held four 3D sonic anemometers (Gill Windmaster Pro) each, at four heights (0.25 m, 1 m, 2 m, 3.5 m), measuring wind speed and direction at 0.1 Hz. The high frequency wind and gas concentration data were combined to measure greenhouse gas concentrations upwind and downwind of the pile, and calculate the flux of CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O from composting, using MMB [11, 24]. The flux ( $\text{g m}^{-2} \text{s}^{-1}$ ) from the source was approximated:

$$\text{Flux} = \frac{1}{L} \int_0^{\infty} \bar{u}_z (\bar{c}_{z,d} - \bar{c}_{z,u}) dz, \quad (1)$$

where  $L$  (m) is the fetch (horizontal distance that air travels over the source),  $\bar{u}_z$  represents mean horizontal wind speed at height  $z$  ( $\text{m s}^{-1}$ ), and  $\bar{c}_{z,d}$  and  $\bar{c}_{z,u}$  are the mean gas concentrations ( $\text{g m}^{-3}$ ) at height  $z$  downwind and upwind of the source. The flux describes the mass of trace gas emitted per unit time, per cross-sectional area of the source (the windrow pile).


### Statistical analyses and emissions calculations

Statistical analyses were performed using the open-source statistical software ‘R.’ Sensor readings were logged every half hour for the duration of the experiment. Two-way, repeated measures ANOVA was used to test the null hypotheses that sensor values did not change by week or position (top, middle, or bottom of pile), and to test variability of weekly physical and chemical properties of the composting material (pH, porosity, N mineralization, nitrification, C:N, moisture).

Wind speed and direction, and gas concentrations were averaged in 8 min blocks, and fluxes were calculated using equation (1), after screening for wind direction. An angle allowance of  $\pm 30$  degrees was used to determine upwind and downwind towers for each time block [9]. Relationships between CH<sub>4</sub> fluxes and environmental variables (temperature, moisture, and O<sub>2</sub>) were examined using breakpoint analyses (R package ‘strucchange’). The number of breakpoints (1, 2 or 3) with the lowest Bayesian Information Criterion (BIC) was selected; breakpoint significance was determined using a structural change test [25].

Emissions factors were estimated by calculating the area under the mean daily flux curves, for each greenhouse gas, using the R package ‘pracma’; we multiplied this integral ( $\text{g m}^{-2}$ ) by the pile volume ( $\text{m}^3$ ) divided by the fetch (m) to estimate the total emissions. For CO<sub>2</sub> fluxes, we additionally perform a mass balance on C, using laboratory analyses of C content in the composting material over the course of the experiment. We compared our emissions factors to those in the literature, by performing a meta-analysis of published studies that measured greenhouse gas emissions from composting.

To estimate net emissions from composting, we created a lifecycle model, using a functional unit of 1 kg of waste (91% manure, 9% green waste by mass) over one year, employing a system boundary beginning at waste generation, and ending at its final application (land or landfill; figure S3). We simplified the model by including only major contributors to greenhouse gas emissions from waste management: avoided emissions, waste processing, and carbon sequestration from land application [26, 27]. To estimate avoided emissions, we assumed that manure would otherwise be managed in an anaerobic lagoon, a common practice in California [2], and yard waste would be used as alternate daily cover in a landfill [28]; emissions values



were taken from Owen and Silver (2015) [2] (manure), and EPA's WARM model (yard waste) [29]. Carbon sequestered from compost application are from Ryals *et al* (2013) [30]. We assumed that 1 kg wet waste produced 0.5 kg compost.

## Results and discussion

### Environmental conditions: temperature, moisture, and oxygen

Temperature varied over time and by position in the compost pile, and followed a pattern consistent with microbial degradation of organic feedstocks [31]. Temperatures rose rapidly initially, then decreased as easily-decomposed substrates were consumed (figure 1(a)). Within that pattern was a weekly oscillation, corresponding to turning events: temperature rose steeply following turning, then declined. The lowest temperatures occurred at the start (day 0), and the highest daily mean temperatures were reached on day 8 ( $74 \pm 0.2$ ,  $71 \pm 0.2$ ,  $68 \pm 0.1$   $^{\circ}\text{C}$ , for top, middle and bottom

positions). Temperature varied significantly between weeks ( $p < 0.05$ ) but not by position in the pile (table S2). Variability of temperature over time is well established in the literature [6, 32]; the composting process is understood as a series of discrete subprocesses (e.g. mesophilic, thermophilic) due to the discontinuity of shifts in temperature and in microbial communities [33]. Variability by location in the pile was observed in another study [34], but was not observed here, possibly due to the high relative lability of manure.

Moisture affects the composting rate and end-product characteristics. Model estimates and field data suggest that wet-weight moisture contents around 50% are best for  $\text{O}_2$  diffusion, water potential, and microbial growth rates during composting [35]. Moisture in the pile varied in response to water consumption and evaporation, and were controlled to keep conditions favorable for decomposition (figure 1(b)). Daily mean moisture varied between  $36\% \pm 0.01\%$  and  $56\% \pm 0.04\%$ . All positions showed moisture increases with pile turning and watering (figure 1(b)). Moisture

differed significantly between weeks ( $p < 0.001$ ), but not by position (table S2).

Oxygen concentrations affect microbial activity, nutrient mineralization, and greenhouse gas dynamics during composting [36]. In this experiment,  $O_2$  concentrations followed an oscillating pattern, with the lowest daily mean values occurring near the beginning and end of the experiment, and highest daily means occurring in the middle (figure 1(c)). Similar patterns have been measured in composting bioreactors [36], while field scale data are highly variable and dependent upon feedstock and management [37]. Daily mean  $O_2$  concentrations ranged from 0.95% to 15% over the 98 d study. The lowest daily mean values were  $1.6 \pm 0.06\%$  (day 65),  $2.3 \pm 0.09\%$  (day 87), and  $0.95 \pm 0.37\%$  (day 0), and the highest daily mean values were  $15 \pm 0.29\%$  (day 19),  $15 \pm 0.09\%$  (day 48), and  $14 \pm 0.09\%$  (day 34), for the bottom, middle and top positions.  $O_2$  level varied significantly by week ( $p < 0.05$ ), and by the interaction between week and position ( $p < 0.05$ ) (table S2). Estimated weekly mean  $O_2$  consumption rates ranged from  $5.3 \pm 0.002 \text{ mg } O_2 \text{ m}^{-2} \text{ s}^{-1}$  (week 1) to  $0.39 \pm 0.0003 \text{ mg } O_2 \text{ m}^{-2} \text{ s}^{-1}$  (week 13). Declining  $O_2$  consumption is consistent with slower microbial decomposition, and declining availability of labile material.

### Biogeochemical characteristics

The average pH values of the composting material ranged from  $7.6 \pm 0.07$  to  $9.3 \pm 0.01$ . pH can be an important predictor of N mineralization rates and denitrification potential. However, N mineralization tends to be most sensitive at pH 7 or lower [38, 39]. pH values did not vary significantly by position, week, or whether samples were taken pre- or post-mixing (table S3a). Porosity varied between  $0.59 \pm 0.02$  and  $0.84 \pm 0.01$ , and varied significantly by week ( $p < 0.001$ ), position ( $p < 0.001$ ), and the interaction between week and position ( $p < 0.05$ ) (table S3b). Porosity provides an index of physical limitations on  $O_2$  diffusion [40, 41]; its value varied over space and time, reflecting feedstock heterogeneity.

The C:N ratio of composting material is an indicator of microbial degradability and the maturity of the composting process [6], and may predict  $CO_2$  and  $N_2O$  emissions [20, 42]. The C:N ratio declined throughout the experiment, mostly due to lowering C concentrations. This decline is consistent with a well-functioning composting process [6]. The C:N ratio varied significantly by week ( $p < 0.001$ ), and by the interaction between position and state of mixing (pre- or post-mixing) ( $p < 0.05$ ) (table S4). The feedstock had initial an C:N of  $26 \pm 0.7$  for all positions, consistent with best management practices (25–30) [6], and final values near 20 ( $21 \pm 0.7$ ,  $21 \pm 0.6$ , and  $19 \pm 0.6$ , for the bottom, middle, and top). Microbes preferentially utilize available forms of C in feedstocks,

and emit  $CO_2$  (as they metabolize aerobically), or  $CO_2$  and  $CH_4$  (anaerobic decomposition) [6].

Potential net nitrification remained near zero throughout the experiment (table S5). Daily mean values ranged between  $-0.20 \pm 0.06$  and  $0.02 \pm 0.01 \text{ mg N g}^{-1}$  compost, and weekly means varied from  $-0.14 \pm 0.04$  to  $0.01 \pm 0.06 \text{ mg N g}^{-1}$ . Potential net nitrification varied significantly with the interaction between position and week, with higher net  $NO_3^-$  consumption in the top position during the second half of the experiment ( $p < 0.05$ ). The low rates of nitrification measured are consistent with high temperature inhibition of nitrification [43, 44].

Potential N mineralization values also bracketed zero, with daily means ranging from  $-0.15 \pm 0.06$  to  $0.34 \pm 0.02 \text{ mg N g}^{-1}$  compost, and weekly means varying from  $-0.11 \pm 0.04$  and  $0.22 \pm 0.08 \text{ mg N g}^{-1}$  compost. Mineralization declined throughout the experiment; values differed significantly by week ( $p < 0.05$ ) (table S5). The highest compost N mineralization rates tend to occur during the thermophilic phase [43, 44].

### Greenhouse gas emissions



Methane fluxes varied over time, with most emissions occurring over short periods early in the composting process. Daily mean  $CH_4$  fluxes ranged from  $17 \mu\text{g } CH_4 \text{ m}^{-2} \text{ s}^{-1}$  to  $4.4 \text{ mg } CH_4 \text{ m}^{-2} \text{ s}^{-1}$ . The highest  $CH_4$  emissions occurred during the first three weeks of composting, during which 75% of total  $CH_4$  was emitted. Half of the total  $CH_4$  emissions occurred in the first 10 d, and 36% occurred in the first week (figure 2(a)). Most fluxes measured were small, with few days contributing the majority of  $CH_4$  emissions. Integrating the daily mean flux curve, the total  $CH_4$  emitted was  $1.7 \pm 0.32 \text{ g } CH_4 \text{ kg}^{-1}$  wet feedstock<sup>2</sup>. This emission factor falls between the median ( $1.4 \text{ g } CH_4 \text{ kg}^{-1}$ ), and mean ( $2.2 \text{ g } CH_4 \text{ kg}^{-1}$ ) literature values found for windrow composting (figure 3, table S1).

Carbon dioxide fluxes were also variable. Daily mean  $CO_2$  fluxes ranged over two orders of magnitude ( $0.3\text{--}30 \text{ mg } CO_2 \text{ m}^{-2} \text{ s}^{-1}$ ) and were an indicator of the C lability of the feedstock. Similar to  $CH_4$ , most  $CO_2$  emissions occurred early, with almost 50% emitted in the first three weeks (figure 2).

Because  $CO_2$  emissions from biological systems are considered to rapidly cycle, these biogenic emissions are usually considered to have no net global warming effect [26, 45–48]. Due to a low sensitivity of the analyzer to  $CO_2$ , the net  $CO_2$  fluxes measured ( $19 \pm 3.7 \text{ g } CO_2 \text{ kg}^{-1}$  wet feedstock<sup>3</sup>) likely present an underestimation of this flux. Utilizing laboratory measurements of C content of the composting material over the course of the experiment to perform a C balance on the composting system, we found an upper

<sup>2</sup>  $3.9 \pm 0.74 \text{ g } CH_4 \text{ kg}^{-1}$  dry.

<sup>3</sup>  $44 \pm 8.6 \text{ g } CO_2 \text{ kg}^{-1}$  dry.



limit on these emissions to be 100 g CO<sub>2</sub> kg<sup>-1</sup>. The upper and lower estimates are within the range of the CO<sub>2</sub> emissions factors found in the literature, as is the high proportion of C loss via CO<sub>2</sub> versus CH<sub>4</sub> (96% to 4% for the upper emissions factor in this study; other studies found even higher proportions lost as CO<sub>2</sub>) [49, 50]. From a greenhouse gas mitigation perspective, it is preferable that the C emitted from composting be emitted as CO<sub>2</sub> rather than as CH<sub>4</sub>.

Nitrous oxide (N<sub>2</sub>O) concentration differences remained below the instrument detection limit (25 ppb + 0.05%), likely reflecting an inability to detect a small signal using micrometeorological methods. The equivalent fluxes of this detection limit (45  $\mu$ g m<sup>-2</sup> s<sup>-1</sup>) were lower than those observed in the literature (table S1; figure S4). Low emissions may be explained by the high C:N measured, above the threshold (25:1) associated with N<sub>2</sub>O emissions [20] for the first four weeks

(and remaining above 20 throughout), during which decomposition is occurring most rapidly. Additionally, as  $\text{NO}_3^-$  is a precursor to  $\text{N}_2\text{O}$ , the low  $\text{NO}_3^-$  concentrations and nitrification rates measured throughout support the lack of  $\text{N}_2\text{O}$  production. Periodic anaerobic conditions and high temperatures likely inhibited nitrification.

Methane was the major contributor to non-biogenic greenhouse gas emissions from composting, and the majority of that  $\text{CH}_4$  was emitted early in the composting process. The timing of emissions suggests that efforts to further decrease direct emissions from composting should focus on increasing  $\text{O}_2$  availability in the early stages, when decomposition and C fluxes (as  $\text{CH}_4$  and  $\text{CO}_2$ ) are at their highest.

We identified 17 studies with 53 estimates of total emissions (emissions factors) from composting [5, 7, 10, 36, 49–61]; the studies spanned different composting processes and methods for emission measurement, feedstocks, and time scales (figure 3; table S1). Most studies used static chamber measurements, which yield periodic measurements of greenhouse gas concentrations at the surface of the composting pile. In the spatially heterogeneous and dynamic composting pile, using static chambers can underestimate fluxes. The measurements reported in the literature varied in duration, from 21 to 150 d. Most studies measured the composting of food and/or green waste; three studies [50, 52, 58] considered manure. The current study used methods to continuously measure whole-pile emissions, exceeded 90 d in duration, and used a high-emitting feedstock (manure); thus, we would expect emissions measured to exceed most of those in the literature. That the resulting emissions factor is near the literature's median value suggests that management may play a larger role than feedstock in predicting greenhouse gas fluxes.

### Emissions patterns and environmental variables

Environmental variables in the composting pile— $\text{O}_2$ , moisture, and temperature—exhibited threshold effects on  $\text{CH}_4$  emissions (figures S5–S7). The break-points (thresholds) for  $\text{CH}_4$  emissions occurred at  $\text{O}_2$  concentrations of 3.3%, 5.7%, and 6.0% ( $p < 0.005$ ); these represent  $\text{O}_2$  levels where the modeled relationship between  $\text{O}_2$  and  $\text{CH}_4$  change [25], with higher emissions occurring below, and lower emissions occurring above those thresholds (figure S5).

In low oxygen environments,  $\text{CH}_4$  emission is controlled by the balance between methanogenesis and methane oxidation. Methanogenesis is favored at low  $\text{O}_2$  concentrations. Methane oxidation rates vary as a function of  $\text{CH}_4$  production,  $\text{O}_2$ , moisture, pH, temperature [62–67], and feedstock chemistry [15, 68] and are limited by diffusive transport [62]. While porosity, a physical control on  $\text{O}_2$  diffusion into the pile, declined over time in our experiment, so did  $\text{CH}_4$  emissions, suggesting that other factors, including

declining temperatures and C availability, played a larger role in controlling emissions than did constraints on  $\text{O}_2$  diffusion. As expected, we observed the highest  $\text{CH}_4$  emissions at lower  $\text{O}_2$  concentrations, coupled with higher moisture, temperatures, and C:N.

Methane fluxes were found to increase more rapidly with temperatures above 65° C ( $p < 0.1$ ) (figure S6). The moisture threshold was found to be 55% ( $p < 0.0001$ );  $\text{CH}_4$  emissions increased more rapidly with moisture as levels exceeded 55% (figure S7). Moisture between 40% and 60% is recommended for composting [6]; higher levels limit  $\text{O}_2$  diffusion into the pile [62], while lower levels preclude microbial activity. The patterns of greenhouse gas emissions from composting can also be sensitive to key biogeochemical characteristics, particularly the relative availability of C and N in the decomposing feedstock. However, here we found no clear predictive (linear or polynomial) relationships between  $\text{CH}_4$  emissions and other chemical or physical characteristics.

Carbon concentrations play an important role in understanding the emissions patterns; the biggest fluxes occur early in composting, and a disproportionate fraction of total emissions occur during the first three weeks, when the feedstock is fresh, and the organic C and N are likely more available to microbes. As composting progresses, the greenhouse gas fluxes decrease as the C and N become more complexed. During the beginning weeks of composting, as microbial degradation speeds up, temperatures and  $\text{O}_2$  consumption were at their highest; these conditions favor methanogenesis.

### Consideration of the larger lifecycle emissions

Total direct greenhouse gas emissions from composting were 57 g  $\text{CO}_2\text{-e kg}^{-1}$  (75–150 g  $\text{CO}_2\text{-e kg}^{-1}$  including biogenic  $\text{CO}_2$ ). Direct emissions are one important, and the most uncertain, element of the lifecycle greenhouse gas emissions from composting. Two other major C fluxes include the avoided emissions—emissions that would have occurred under ‘business as usual’ waste management—and enhanced C sequestration from land application of compost. Using a system boundary that includes these three processes (figure S3), avoided emissions from the landfill (for green waste) and anaerobic lagoons (for manure) outweigh the direct emissions from composting (table 1). The net lifecycle emissions from composting manure and green waste and applying the compost to grasslands are negative (–690 g  $\text{CO}_2\text{-e kg}^{-1}$  excluding biogenic  $\text{CO}_2$ ; between –590 and –670 g  $\text{CO}_2\text{-e kg}^{-1}$  including the directly measured biogenic  $\text{CO}_2$  fluxes)<sup>4</sup>, meaning that emissions are avoided on net. If all of California's dairy manure that is currently managed via anaerobic lagoon (7.4 MMT  $\text{yr}^{-1}$ ) [69] were instead composted, approximately 5MMT  $\text{CO}_2\text{-e yr}^{-1}$

<sup>4</sup> –1.6 kg  $\text{CO}_2\text{-e kg}^{-1}$  dry (–1.4 to –1.5  $\text{CO}_2\text{-e kg}^{-1}$  dry including biogenic  $\text{CO}_2$ ).



**Table 1.** Net lifecycle greenhouse gas emissions from composting green waste and manure,  $\text{g CO}_2\text{-e kg}^{-1}$  feedstock (wet). Net emissions are a sum of direct emissions, C sequestered, and a weighted sum of avoided emissions from anaerobic lagoons (for manure, 91% of mass) and landfills (for green waste, 9% of mass).

Emission source	Value
Total direct emissions (no biogenic $\text{CO}_2$ )	+5.6 E + 01
Avoided, anaerobic lagoon (manure)	−8.1 E + 02
Avoided, landfill (green waste)	−3.3 E + 01
Carbon sequestered from land application	−2.8 E − 01
Net emissions	−6.9 E + 02

would be avoided; this is equivalent to 15% of California's agricultural greenhouse gas emissions.

The biggest greenhouse gas benefit from composting is the elimination of anaerobic storage. Emissions avoided from anaerobic storage were greater than the direct emissions from composting. Though annual C sequestration from compost application is smaller in magnitude than direct and avoided emissions, its effects are potentially long-lasting. Avoided and direct emissions are a one-time phenomenon, while increased C sequestration from a single compost application can persist for several years [70]. Thus, while we modeled net C implications over one year (table 1), a longer time frame would increase the relative benefits of compost application.

Our results highlight the potential to effectively measure and minimize greenhouse gas emissions from commercial-scale composting. Managing composting piles to minimize methanogenesis—by maintaining sufficient  $\text{O}_2$  concentrations through aeration and bulking, and focusing on the first three weeks of decomposition when temperatures and decomposition rates are high—could potentially reduce direct greenhouse gas emissions. Low emissions from composting contribute to the climate change mitigation benefit of diverting high-emitting waste streams to compost and land applying for C sequestration.

## Acknowledgments

In addition to the funding sources, the authors would also like to especially thank John Wick, Tyler Anthony, Heather Dang, Summer Ahmed, and Joseph Verfaillie for their support of this work, as well as Picarro Instruments, Gill Instruments, and NovaLynx Corporation. We would also like to thank the three anonymous reviewers for their thoughtful and valuable feedback.

## Data availability statement

The data that support the findings of this study are available from the corresponding author upon reasonable request.

## Author contributions

The manuscript was written through contributions of both authors. Both authors have given approval to the final version of the manuscript.

## Funding sources

This work was supported by funding from California's 4th Climate Change Assessment, through the Berkeley Energy and Climate Initiative, and the Rathmann Family Foundation. Additional support was provided to WLS from the 11th Hour Foundation.

## ORCID iDs

Sintana E Vergara  <https://orcid.org/0000-0002-1322-7568>

## References

- [1] Bogner J, Abdelrafie Ahmed M, Diaz C, Faaij A, Gao Q, Hashimoto S, Mareckova K, Pipatti R and Zhang T 2007 *Waste Management. In Climate Change 2007: Mitigation. Contribution of Working Group III to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change* (Cambridge and New York, NY: Cambridge University Press)
- [2] Owen J J and Silver W L 2015 Greenhouse gas emissions from dairy manure management: a review of field-based studies *Glob. Change Biol.* **21** 550–65
- [3] King F H 1911 *Farmers of Forty Centuries; or, Permanent Agriculture in China, Korea and Japan* (Madison, WI: F H King)
- [4] Myhre G *et al* 2013 Anthropogenic and natural radiative forcing *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* (Cambridge and New York, NY: Cambridge University Press) pp 659–740
- [5] Hellmann B, Zelles L, Palojarvi A and Bai Q 1997 Emission of climate-relevant trace gases and succession of microbial communities during open-windrow composting *Appl. Environ. Microbiol.* **63** 1011–8
- [6] Diaz L F and Savage G M 2007 Factors that affect the process *Compost Science and Technology (Waste Management Series)* ed L F Diaz *et al* vol 8 (Amsterdam: Elsevier) pp 49–65
- [7] Amlinger F, Peyr S and Cuhls C 2008 Green house gas emissions from composting and mechanical biological treatment *Waste Manage. Res.* **26** 47–60
- [8] Sommer S G, McGinn S M, Hao X and Larney F J 2004 Techniques for measuring gas emissions from a composting stockpile of cattle manure *Atmos. Environ.* **38** 4643–52
- [9] Wagner-Riddle C, Park K-H and Thurtell G W 2006 A micrometeorological mass balance approach for greenhouse gas flux measurements from stored animal manure *Agric. For. Meteorol.* **136** 175–87
- [10] Zhu-Barker X, Bailey S K, Paw U K T, Burger M and Horwath W R 2017 Greenhouse gas emissions from green waste composting windrow *Waste Manage.* **59** 70–9
- [11] Harper L A, Denmead O T and Flesch T K 2011 Micrometeorological techniques for measurement of enteric greenhouse gas emissions *Animal Feed Sci. Technol.* **166–167** 227–39
- [12] Kent E 2010 Micrometeorological measurements of GHG emissions from composting yard waste *Master's Thesis* University of California Davis
- [13] Chapuis-Lardy L, Wrage N, Metay A, Chotte J-L and Bernoux M 2007 Soils, a sink for  $\text{N}_2\text{O}$ ? A review *Glob. Change Biol.* **13** 1–17

- [14] Bridgham S D and Richardson C J 1992 Mechanisms controlling soil respiration (CO<sub>2</sub> and CH<sub>4</sub>) in southern peatlands *Soil Biol. Biochem.* **24** 1089–99
- [15] Luo W H, Yuan J, Luo Y M, Li G X, Nghiem L D and Price W E 2014 Effects of mixing and covering with mature compost on gaseous emissions during composting *Chemosphere* **117** 14–9
- [16] Conrad R 1996 Soil Microorganisms as controllers of atmospheric trace gases (H<sub>2</sub>, CO, CH<sub>4</sub>, OCS, N<sub>2</sub>O, and NO) *Microbiol. Rev.* **60** 609–40
- [17] Treat C C, Wollheim W M, Varner R K, Grandy A S, Talbot J and Frohling S 2014 Temperature and peat type control CO<sub>2</sub> and CH<sub>4</sub> production in alaskan permafrost peats *Glob. Change Biol.* **20** 2674–86
- [18] Olefeldt D, Turetsky M R, Crill P M and McGuire A D 2013 Environmental and physical controls on northern terrestrial methane emissions across permafrost zones *Glob. Change Biol.* **19** 589–603
- [19] Firestone M K and Davidson E A 1989 Microbiological basis of NO and N<sub>2</sub>O production and consumption in soil *Exchange of Trace Gases between Terrestrial Ecosystems and the Atmosphere* ed M O Andreae and D S Schimel (Chichester: Wiley) pp 7–21
- [20] Klemetsson L, Arnold K V, Weslien P and Gundersen P 2005 Soil CN ratio as a scalar parameter to predict nitrous oxide emissions *Glob. Change Biol.* **11** 1142–7
- [21] McLean E O 1982 Soil PH and lime requirement *Methods of soil analysis, Part 2* ed A L Page *et al* (Madison, WI: American Society of Agronomy) pp 199–224
- [22] Danielson R E and Sutherland P L 1986 Porosity *Methods of Soil Analysis Part 1. Physical and Mineralogical Methods, Agronomy Monograph no. 9* (Madison, WI: Soil Science Society of America) pp 443–61
- [23] Hart S C, Stark J M, Davidson E A and Firestone M K 1994 Nitrogen mineralization, immobilization, and nitrification *Methods of Soil Analysis: Part 2—Microbiological and Biochemical Properties* (Madison, WI: Soil Science Society of America) pp 985–1018
- [24] Denmead O T 2008 Approaches to measuring fluxes of methane and nitrous oxide between landscapes and the atmosphere *Plant Soil* **309** 5–24
- [25] Zeileis A 2006 Implementing a class of structural change tests: an econometric computing approach *Comput. Stat. Data Anal.* **50** 2987–3008
- [26] Vergara S E, Damgaard A and Horvath A 2011 Boundaries matter: greenhouse gas emission reductions from alternative waste treatment strategies for california’s municipal solid waste *Resour. Conserv. Recycle* **57** 87–97
- [27] DeLonge M S, Ryals R and Silver W L 2013 A lifecycle model to evaluate carbon sequestration potential and greenhouse gas dynamics of managed grasslands *Ecosystems* **16** 962–79
- [28] Cal Recycle 2016 *State of Recycling in California: Updated 2016 DRRR-2016-01554* Sacramento CA: California Department of Resources Recycling and Recovery State of Recycling in California: Updated 2016 DRRR-2016-01554
- [29] United States Environmental Protection Agency 2019 Waste Reduction Model (WARM), Washington DC (<https://www.epa.gov/warm>)
- [30] Ryals R and Silver W L 2013 Effects of organic matter amendments on net primary productivity and greenhouse gas emissions in annual grasslands *Ecol. Appl.* **23** 46–59
- [31] Insam H, Amor K, Renner M and Crepez C 1996 Changes in functional abilities of the microbial community during composting of manure *Microb. Ecol.* **31** 77–87
- [32] Finstein M S and Morris M L 1975 Microbiology of municipal solid waste composting 11 paper of the journal series, new jersey agricultural experiment station *Advances in Applied Microbiology* ed D Perlman (New York: Academic) vol 19, pp 113–51
- [33] Insam H and de Bertoldi M 2007 Microbiology of the composting process *Compost Science and Technology (Waste Management Series)* ed L F Diaz *et al* vol 8 (Amsterdam: Elsevier) pp 25–48
- [34] Fernandes L, Zhan W, Patni N K and Jui P Y 1994 Temperature distribution and variation in passively aerated static compost piles *Bioresour. Technol.* **48** 257–63
- [35] Richard T L, Hamelers H V M, (Bert), Veeken A and Silva T 2002 Moisture relationships in composting processes *Compost Sci. Util.* **10** 286–302
- [36] Puyuelo B, Gea T and Sánchez A 2014 GHG emissions during the high-rate production of compost using standard and advanced aeration strategies *Chemosphere* **109** 64–70
- [37] Michel F C, Huang J F, Forney L J and Reddy C A 1996 Field scale study of the effect of pile size, turning regime and leaf to grass mix ratio on the composting of yard trimmings *The Science of Composting* ed M de Bertoldi *et al* (Dordrecht: Springer Netherlands) pp 577–84
- [38] Curtin D, Campbell C A and Jalil A 1998 Effects of acidity on mineralization: PH-dependence of organic matter mineralization in weakly acidic soils *Soil Biol. Biochem.* **30** 57–64
- [39] Samad M S, Bakken L R, Nadeem S, Clough T J, Klein C A M, de; Richards K G, Lanigan G J and Morales S E 2016 High-resolution denitrification kinetics in pasture soils link N<sub>2</sub>O emissions to PH, and denitrification to C mineralization *PLoS One* **11** e0151713
- [40] Ruggieri L, Gea T, Artola A and Sánchez A 2009 Air filled porosity measurements by air pycnometry in the composting process: a review and a correlation analysis *Bioresour. Technol.* **100** 2655–66
- [41] Stepniewski W, GLińSKI J and Ball B C 1994 Effects of compaction on soil aeration properties *Developments in Agricultural Engineering* ed B D Soane and C van Ouwerkerk *Soil Compaction in Crop Production* (Amsterdam: Elsevier) vol 11, pp 167–89
- [42] Huang Y, Zou J, Zheng X, Wang Y and Xu X 2004 Nitrous oxide emissions as influenced by amendment of plant residues with different C:N ratios *Soil Biol. Biochem.* **36** 973–81
- [43] Bernal M P, Albuquerque J A and Moral R 2009 Composting of animal manures and chemical criteria for compost maturity assessment. a review *Bioresour. Technol.* **100** 5444–53
- [44] Tiquia S M 2002 Microbial transformation of nitrogen during composting *Microbiology of Composting* ed H Insam *et al* (Berlin: Springer) 237–45
- [45] Rabl A, Benoist A, Dron D, Peuportier B, Spadaro J V and Zoughaib A 2007 How to account for CO<sub>2</sub> emissions from biomass in an LCA *Int. J. Life Cycle Assess.* **12** 281
- [46] Gentil E, Christensen T H and Aoustin E 2009 Greenhouse gas accounting and waste management *Waste Manage. Res. J. Int. Solid Wastes Public Clean. Assoc. ISWA* **27** 696–706
- [47] Christensen T H, Gentil E, Boldrin A, Larsen A W, Weidema B P and Hauschild M C 2009 Balance, carbon dioxide emissions and global warming potentials in LCA-modelling of waste management systems *Waste Manage. Res.* **27** 707–15
- [48] Barton J R, Issaia I and Stentiford E I 2008 Carbon—making the right choice for waste management in developing countries *Waste Manage.* **28** 690–8
- [49] Boldrin A, Andersen J K, Møller J, Christensen T H and Favoino E 2009 Composting and compost utilization: accounting of greenhouse gases and global warming contributions *Waste Manage. Res.* **27** 800–12
- [50] Andersen J K, Boldrin A, Christensen T H and Scheutz C 2010 Greenhouse gas emissions from home composting of organic household waste *Waste Manage.* **30** 2475–82
- [51] Adhikari B K, Trémier A, Barrington S, Martinez J and Daumoin M 2013 Gas emissions as influenced by home composting system configuration *J. Environ. Manage.* **116** 163–71
- [52] Amon B, Amon T and Boxberger J 2001 Alt, Ch. emissions of NH<sub>3</sub>, N<sub>2</sub>O and CH<sub>4</sub> from dairy cows housed in a farmyard manure tying stall (housing, manure storage, manure spreading) *Nutr. Cycl. Agroecosyst.* **60** 103–13
- [53] California Air Resources Board (ARB) 2017 *Method for estimating greenhouse gas emission reductions from diversion of organic waste from landfills to compost facilities* Industrial

- Strategies Division, Transportation and Toxics Division, California Air Resources Board (<https://ww3.arb.ca.gov/cc/waste/cerffinal.pdf>)
- [54] Colón J, Cadena E, Pognani M, Barrera R, Sánchez A, Font X and Artola A 2012 Determination of the energy and environmental burdens associated with the biological treatment of source-separated municipal solid wastes *Energy Environ. Sci.* **5** 5731–41
- [55] Cuhls C, Mahl B and Clemens J 2015 *Ermittlung der Emissionssituation bei der Verwertung von Bioabfällen* (Dessau-Roßlau: Umweltbundesamt)
- [56] de Groot M 2010 *Update of emission factors for N<sub>2</sub>O and CH<sub>4</sub> for composting, anaerobic digestion and waste incineration C9910-01-001* DHV B.V., The Netherlands p 33 ([https://www.rvo.nl/sites/default/files/2013/10/DHV2010%20-%20Update%20emission%20factors%20N<sub>2</sub>O%20and%20CH<sub>4</sub>%20for%20Waste.pdf](https://www.rvo.nl/sites/default/files/2013/10/DHV2010%20-%20Update%20emission%20factors%20N2O%20and%20CH4%20for%20Waste.pdf))
- [57] Ermolaev E, Jarvis Å, Sundberg C, Smårs S, Pell M and Jönsson H 2015 Nitrous oxide and methane emissions from food waste composting at different temperatures *Waste Manage.* **46** 113–9
- [58] He Y, Inamori Y, Mizuochi M, Kong H, Iwami N and Sun T 2001 Nitrous oxide emissions from aerated composting of organic waste *Environ. Sci. Technol.* **35** 2347–51
- [59] Hellebrand H J 1998 Emission of nitrous oxide and other trace gases during composting of grass and green waste *J. Agric. Eng. Res.* **69** 365–75
- [60] San Joaquin Valley Air Pollution Control District 2013 *Greenwaste compost site emissions reductions from solar-powered aeration and biofilter layer* San Joaquin Valley Technology Advancement Program p132 (<https://www.o2compost.com/Userfiles/PDF/VOC-Emissions-Report.pdf>)
- [61] South Coast Air Quality Management District 2001 *Remote sensing tests for ammonia and volatile organic compound emissions from a greenwaste composting operation* Monitoring and source test engineering branch, Monitoring and analysis, South Coast Air Quality Management District p 11 ([https://ww3.arb.ca.gov/cc/compost/documents/inlandempire2\\_source.pdf](https://ww3.arb.ca.gov/cc/compost/documents/inlandempire2_source.pdf))
- [62] King G 1997 Responses of atmospheric methane consumption by soils to global climate change *Glob. Change Biol.* **3** 351–62
- [63] Teh Y A, Silver W L and Conrad M E 2005 Oxygen effects on methane production and oxidation in humid tropical forest soils *Glob. Change Biol.* **11** 1283–97
- [64] Teh Y A, Dubinsky E A, Silver W L and Carlson C M 2008 Suppression of Methanogenesis by dissimilatory Fe(III)-reducing bacteria in tropical rain forest soils: implications for ecosystem methane flux *Glob. Change Biol.* **14** 413–22
- [65] Teh Y A and Silver W L 2006 Effects of soil structure destruction on methane production and carbon partitioning between methanogenic pathways in tropical rain forest soils *J. Geophys. Res. Biogeosci.* **111** 1–8
- [66] Sullivan B W, Selmants P C and Hart S C 2014 What is the relationship between soil methane oxidation and other C compounds? *Glob. Change Biol.* **20** 2381–2
- [67] McNicol G and Silver W L 2014 Separate effects of flooding and anaerobiosis on soil greenhouse gas emissions and redox sensitive biogeochemistry *J. Geophys. Res. Biogeosci.* **119** 557–66
- [68] Sonoki T, Furukawa T, Jindo K, Suto K, Aoyama M and Sánchez-Monedero M Á 2013 Influence of biochar addition on methane metabolism during thermophilic phase of composting *J. Basic Microbiol.* **53** 617–21
- [69] California Air Resources Board (ARB) 2014 *Annex 3B. Manure Management (IPCC 3A2) to the Technical Support Document for the 2000-2012 California's Greenhouse Gas Emissions Inventory* State of California Air Resources Board Air Quality Planning and Science Division ([https://ww3.arb.ca.gov/cc/inventory/pubs/reports/2000\\_2012/ghg\\_inventory\\_00-12\\_technical\\_support\\_document.pdf](https://ww3.arb.ca.gov/cc/inventory/pubs/reports/2000_2012/ghg_inventory_00-12_technical_support_document.pdf))
- [70] Ryals R, Hartman M D, Parton W J, DeLonge M S and Silver W L 2015 Long-term climate change mitigation potential with organic matter management on grasslands *Ecol. Appl.* **25** 531–45